

Washington
2008 School Health Profiles Report
UNDERSTANDING YOUR PROFILES RESULTS

This section is designed to assist you with understanding the different type of results presented in the School Health Profiles (Profiles) report and with how to interpret the findings. The following topics are included:

- Understanding Sample Descriptions, Response Rates and Weighting Procedures
- Understanding Your Summary of Principal and Teacher Results
- Understanding Your Principal Teacher Results Tables
- Understanding Your Principal and Teacher Results Charts
- Understanding Your SLIMs Results

Understanding Sample Descriptions, Response Rates and Weighting Procedures

Being able to describe the population the results represent and provide support for generalizing the results to the population is important when presenting the results. Your ability to generalize the results of your Profiles is determined by whether your data are unweighted or weighted. If your survey response rate is less than 70%, your data are unweighted. If it is 70% or greater, you have used a scientifically selected sample, and you provided all necessary documentation, your data have been weighted.

If the data are unweighted, the data in the tables and charts refer only to those schools whose principals or teachers actually participated in the survey. For example, if you have unweighted data and 30% of schools in the sample report they require health education, you can report that 30% of the schools that participated in the survey require health education.

If the data are weighted, the percentages refer to all schools in that category within your district or state. For example, if the data are weighted and 30% of the schools in the high school category report they require health education, you can report that 30% of high schools in your state or district require health education.

The Principal Survey and the Teacher Survey are weighted separately. This report has a sample description section for the both the Principal and Teacher Survey. It describes the type of schools surveyed (e.g., all regular secondary public schools having at least one of grades 6 through 12) and the method used to select the sample of schools.

The number of principals/teachers in the sample, the number of eligible principals/teachers who returned the questionnaires, the response rate, and how the weights are calculated are also provided. This report

Understanding Your Summary of Principal Teacher Results

The Summary presents selected results (percents and confidence intervals) from both the Principal and Teacher surveys. This summary provides quick reference for several key results in one table, by topic, and may serve as a ready-to-use fact sheet about your Profiles survey.

Megatropolis 2008 School Health Profiles Report Summary of Principal and Lead Health Education Teacher Survey Results		
	Percent	95% Confidence Interval
HEALTH EDUCATION		
Percent of schools that require students to take 2 or more health education courses	54	49 - 59
Among schools that require a health education course, percent that require students who fail the course to repeat it	81	78 - 84
Percent of schools in which the lead health education teacher had professional preparation in health education or in health and physical education combined	19	16 - 22
Percent of schools in which the lead health education teacher is certified, licensed, or endorsed by the state to teach health education in middle school or high school	28	24 - 33
PHYSICAL EDUCATION AND PHYSICAL ACTIVITY		
Among schools that require physical education, percent of schools in which students can not be exempted from taking required physical education for one grading period or longer*	35	31 - 39

Example: 54% of schools in your state require students to take 2 or more health education courses.

The 95% confidence interval is explained on page 6 of this booklet.)

Understanding Your Principal and Teacher Results Tables

The report provides Tables of all your results. The example table below will help illustrate the format.

Megatropolis 2008 School Health Profiles Report Weighted Principal Survey Results			
14. Percentage of schools that require all staff who teach physical education to be certified, licensed, or endorsed by the state in physical education.			
	N	Percent	95% Confidence Interval
a. Yes			
High Schools	27	92	82 - 97
Middle Schools	24	67	58 - 75
Junior/Senior High Schools	103	20	15 - 25
All Schools	154	39	36 - 43
b. No			
High Schools	27	8	3 - 18
Middle Schools	24	33	25 - 42
Junior/Senior High Schools	103	73	68 - 78
All Schools	154	56	52 - 60
c. Not Applicable			
High Schools	27	0	NA
Middle Schools	24	0	NA
Junior/Senior High Schools	103	7	5 - 10
All Schools	154	5	3 - 7

Title

The title of each table tells whether the data describe information from school principals or lead health education teachers, and whether the data are weighted or unweighted.

The question number and a summary of the question appear under the title. Response options for the question or subparts of the question are listed on the far left of the table. In some tables, question numbers are followed by the letter N (e.g., 21N). The data contained in these tables were derived from combining two or more response options to a single question or two or more questions or question subparts. Where this occurs, the response options, questions, or question subparts used in creating the combined variable are listed in a footnote.

The results are provided for **All Schools** regardless of weight status and each table also lists responses for three categories of schools:

- High schools, defined as those with a low grade of 9 or higher and a high grade of 10 or higher;
- Middle schools, defined as those with a high grade of 9 or lower; and
- Junior/senior high schools, defined as those with a low grade of 8 or less and a high grade of 10 or higher.

“N”

In addition to the column containing the response options, each table has two or three other columns. The **N** column shows the total unweighted number of observations for each category of school (high school, middle school, and junior/senior high school) and all schools. Stated another way, N is the number of schools in each category and overall that provided any response to the question or subparts of the question. For example, in the above table, the principals in 27 high schools answered question 14.

Percent

The **Percent** column shows the percentage of schools in each category and overall that responded to the question as described in the question summary at the top of each table. If the data are weighted, this is a weighted percentage; if the data are unweighted, it is an unweighted percentage. In the example table above, 92% of high schools in the state require all staff who teach physical education to be certified, licensed, or endorsed by the state in physical education.

95% Confidence Intervals

A sample of schools were selected to take the survey so the results are weighted, a fourth column shows the **95% Confidence Interval** for the percentage reported. The 95% confidence interval provides the range of values within which the "true" percentage lies. A 95% confidence interval means that if the survey were repeated many times, the "true" value would fall within the interval 95% of the time.

A census of all eligible schools was used for the Washington Supplemental Survey, so the results are unweighted. The response rate for the supplements were less than 70% so 95% confidence intervals were also provided.

When the confidence interval is relatively narrow, you have a more precise indication of the requirement for staff who teach physical education to be certified, licensed, or endorsed in physical education. Wider confidence intervals diminish the ability to report results with precision. In the example table above, the confidence interval is 82 to 97. This means you can be 95% confident that the "true" percentage of schools could be as low as 82% or as high as 97%.

Determining Significant Differences

If you are interested in comparing two percentages, confidence intervals can also be used for comparisons.

If the confidence intervals of two groups you are interested in comparing do NOT overlap, the results are considered significantly different.

- Using the results from the above example table, we can compare the percentage of high schools 92% (82 -97) to the percentage of middle schools 67% (58 – 75) that have a requirement for certification, licensure, or endorsement for physical education teachers.
- The two confidence intervals do not overlap – therefore, the percentage of high schools that require all staff who teach physical education to be certified, licensed, or endorsed by the state in physical education is significantly higher than the percentage of middle schools with the same requirement.

If at least one of the confidence intervals overlaps one of the percent estimates the two results are Not considered significantly different.


If the confidence intervals overlap, but the overlap does not include one the percent estimates, then the differences is inconclusive.

- If your confidence intervals give you an inconclusive results, you can determine statistical significance by using an online tool is available at <http://www.hys.wa.gov/Reporting/Default.aspx>

Understanding Your Principal and Teacher Results Charts

For each table, a bar chart presenting the same results graphically is provided. Across the top of each bar chart is the title indicating whether the data are from school principals or lead health education teachers and whether the data in the chart are weighted or unweighted. The question number and summary of the question appear below the title. The response options or subparts of the question are listed down the left margin of the chart. Each item in the chart contains four bars: one for each of three categories of schools (high school, middle school, and junior/senior high school) and one for all schools combined (all schools). Each bar represents the percentage of schools in each category and overall that responded to the question as described in the question summary at the top of each chart. The percent is to the right of the bar.

A sample bar chart appears below. The charts are useful for creating presentations or adding to your own Profiles report.



Understanding Your SLIMs Results

Sites who received funding for Priorities 2, 3, or 4 of the CDC/DASH cooperative agreement (DP08-801) are required to “document the impact of their program activities by monitoring the percentage of schools in their jurisdiction that are implementing specific, effective...policies and practices.” The results from your 2008 Profiles survey can be used to help you choose which School-Level Impact Measures (SLIMs) you will focus on and future Profiles survey results can be used to monitor your progress.

In this section of the report, tables describe each SLIM and provide the percentage of schools meeting the requirements for the SLIM. The title indicates which set of SLIMs are provided (the example below presents the results for the human immunodeficiency virus (HIV) prevention SLIMs). Below the title are two columns. The column on the left identifies the SLIM. The column on the right provides the results.

Megatropolis	
School-Level Impact Measures (SLIMs)	
Part I: Human Immunodeficiency Virus (HIV) Prevention SLIMs	
School-Level Impact Measure (SLIM)	Percentage of Schools Meeting SLIM
HIV 1. The percentage of schools that address all of the following in a required course taught during grades 6, 7, or 8: <ul style="list-style-type: none"> ▪ The differences between HIV and AIDS. ▪ How HIV and other STD are transmitted. ▪ How HIV and other STD are diagnosed and treated. ▪ Health consequences of HIV, other STD, and pregnancy. ▪ The benefits of being sexually abstinent. ▪ How to prevent HIV, other STD, and pregnancy. ▪ How to access valid and reliable health information, products, and services related to HIV, other STD, and pregnancy. ▪ The influences of media, family, and social and cultural norms on sexual behavior. ▪ Communication and negotiation skills related to eliminating or reducing risk for HIV, other STD, and pregnancy. ▪ Goal setting and decision making skills related to eliminating or reducing risk for HIV, other STD, and pregnancy. ▪ Compassion for persons living with HIV or AIDS. 	32
HIV 2. The percentage of schools that address all of the following in a required course taught during grades 9, 10, 11, or 12: <ul style="list-style-type: none"> ▪ The relationship among HIV, other STD, and pregnancy. 	48

In the example above, 32% of schools in the jurisdiction met all requirements for the HIV 1 SLIM.

Please contact your CDC/DASH Project Officer for more information on how to choose the appropriate SLIMs and how to interpret the results as part of your cooperative agreement.

Washington
2008 School Health Profiles Report
USING YOUR PROFILES RESULTS

This section is designed to assist you with presenting your Profiles results effectively. It includes guidelines for planning and developing reports, choosing the method for reporting, and using effective graphics. It also describes additional data sources you can use to supplement your Profiles results. The following topics are included:

- Planning and Developing Effective Reports
- Using Effective Graphics
- Checklist for Effective Graphics

Effective reporting of your Profiles results enables you to provide a broad audience with factual information on school health policies and programs in the following areas: health education; physical education and activity; health services; healthy and safe school environment; and family and community involvement. Concrete recommendations can be made to education agencies, public health officials, parents, and those who assist in the development of your health education programs. The results contained in your report can be used to provide the support for those recommendations.

Reporting your results to the appropriate audiences in an effective and timely manner can:

- Increase commitments to support school health policies and programs;
- Help you make concrete, data-supported recommendations for school health policies and programs in your district or state;
- Enable you to respond more easily and effectively to public and media requests for information about school health policies and programs; and
- Encourage increased participation in future years.

Planning and Developing Effective Reports

When planning your report, consider the following:

- audience,
- content,
- style and format, and
- method of reporting.

Determining your **audience** is the first step in planning and developing your report or reports. A primary audience for your Profiles is CDC, but legislators, school board members, and district/state administrators are other audiences for your Profiles. Education department officials and program staff, teachers, trainers, parents, and the community may also be interested in your results.

Your audience will determine the **content** of the report. Vary the materials you use for reporting to different audiences and consider the following for each group:

- existing levels of knowledge;
- key concerns and issues;
- method of presentation most likely to draw that group's attention; and
- types of information most likely to motivate action.

Emphasize those aspects of your Profiles data that are most interesting to each audience. Focus on the most important points you want to make instead of overwhelming your audience with details. Provide only the level of detail the audience needs or has requested.

Your results will be more meaningful if presented with other relevant data. You may want to compare your district or state data to data that reflect general national results. Other potential sources of data include:

- * Previous Profiles results;
- * The School Health Policies and Programs Study 2006;
- * Your district or state Youth Risk Behavior Survey (YRBS), Youth Tobacco Survey (YTS), and other student surveys;
- * The national school-based YRBS and the national YTS;
- * Your state Behavioral Risk Factor Surveillance System (BRFSS);
- * Survey results from other states and cities and national organizations;
- * Health outcome data;
- * Item-by-item rationale for the principal and teacher questionnaires for the Profiles (included in your Profiles report binder); and
- * National health statistics from federal agencies.

Examples of combining the Youth Risk Behavior Survey (YRBS) results and the Profiles results can be found on the CDC/DASH website (<http://www.cdc.gov/healthyyouth/profiles/index.htm>). If current YRBS and Profiles results were available, three fact sheets were created for sites on the following topics – Childhood Overweight, HIV, Tobacco Use, and Violence.

Consider the **style and format**, as well as the content of the product you will target to each audience you want to reach. Organize your findings in a way that clarifies implications for each particular audience. Reports that contain visual images are more powerful than those with just written text.

Once you have selected the findings and determined a style and format that will have the most impact, the final step is to develop the appropriate **method** for dissemination. Keep in mind, using more than one method has a better chance of reaching and persuading a broad audience.

Examples of methods that can be used include:

Executive summary. A two- or three-page executive summary should include all the relevant information generally needed for the reader to become informed about the subject. Use this reporting mechanism to make recommendations for change based on the reported information.

Comprehensive report. Include many or all findings and details from your Profiles in a comprehensive report. Use bullets, boxes, and graphics to emphasize what you want the reader to remember. Include your executive summary as an introduction. Organize the report by topic and include principal data and teacher data in each section as appropriate.

Newsletter. Use a newsletter to report information specifically addressed to certain groups of people, such as teachers, parents, or other professional or community groups. Contribute to existing newsletters or develop your own. Publishing in the state/district principal and teacher newsletters is a way of gaining support for future surveys.

Fact sheets and brochures. A single-page fact sheet or threefold brochure should focus on three or four key results. Fact sheets also might contain some information about your YRBS results. Include your project name, address, and telephone number. Fact sheets can be used easily to disseminate information widely.

Visual presentation. In addition to written reports, you may want visual presentations to report your results before an audience. Consider overhead transparencies, slides, or a PowerPoint presentation depicting the most important findings from your Profiles. Include text interspersed with graphs that focus on a single finding.

Website. You may want to include your findings on an existing website or create a website so you can share your Profiles data more easily and with many interested parties.

Using Effective Graphics

In reporting statistical data, graphic representation can be extremely useful in displaying results in an easy-to-understand manner. Graphics are charts, graphs, and other visual forms for presenting information. Graphic presentation of data is a powerful tool when effectively used. Graphic enhancements are often the sparks that bring life, attention, and interest to a report or presentation. Graphic images help demonstrate group differences and aid in the explanation of survey findings.

The remainder of this booklet has been developed to help you prepare accurate and effective graphics. It focuses primarily on graphics used in presentation but the same guidelines can be used when including graphics in any report method. The guidelines are not intended to constrain creativity, but rather to encourage and support accuracy and consistency in the display of information. Your Profiles report CD-ROM contains graphs for all of your questions in a PowerPoint presentation format. If you want to add additional “slides” or modify this presentation, you can make these changes yourself or find out about services available in your education or health agency.

Planning Your Graphic Presentation

The first step to preparing effective graphic presentations is to ensure that they have a clear purpose. Think about what you are trying to say with the graphic. Keep your message simple and straightforward. Remember that your graphic presentation should highlight your major findings.

Graphic presentations provide an opportunity for you to acquaint various audiences with your program. You must know your audience members so you can design a presentation to best fit their needs. For example, knowing whether your graphics will be viewed by policy makers, such as district superintendents, or by parent groups will help you decide how to present your results.

A graphic's primary function is to inform. This can best be done when data are presented clearly and simply. Simple graphics that are easy to understand will communicate your survey findings much more effectively than tables of raw data. Ideally, your graphics should be both accurate and visually appealing.

Graphics within a presentation should have a consistent style and format. Although many type or font styles are available, using too many different styles can add an inconsistent, cluttered, unprofessional look to an otherwise clean and simple presentation. If you add “slides” to your Profiles report presentation, limit your choices to one or two fonts, and use boldface or italics for emphasis.

Another key factor to consider is the amount of information to convey in a single graphic. Too much information makes a graphic difficult to comprehend, which in turn detracts from your ability to demonstrate important programmatic needs. A series of simple graphics may be far more effective than a single complicated graph. However, be careful not to summarize the information to the point that it misrepresents the actual data.

Keeping presentation graphics as simple as possible forces you to interpret and discuss them in a conversational tone rather than reading them verbatim to your audience. Reading your slides and overhead transparencies is boring for both you and your audience. Your graphics should contain the framework rather than all the details of your presentation.

Several types of charts can be used to display your data. Choose the one that will best highlight the point you want to make.

Text Charts

Use text charts to introduce nonnumeric data in a presentation, for example, to introduce or summarize your findings. Text charts should be short and precise in meaning, using the minimum number of short keywords needed to convey your message. Keep lines short by highlighting only the main idea. Limit text charts to 8 lines, with no more than 8 to 10 words on a line. Paraphrase rather than use complete sentences. Use initial capital letters and lowercase (as in the example shown) for the rest of the text. **USING ALL UPPERCASE LETTERS MAKES TEXT DIFFICULT TO READ.**

Avoid jargon. Be careful when using abbreviations or acronyms. For example, be sure your audience knows that Profiles stands for School Health Profiles.



2008 School Health Profiles Survey

School policies and programs assessed by School Health Profiles (Profiles):

- Health education
- Physical education and activity
- Health services
- Healthy and safe school environment
- Family and community involvement


Use bulleted lists to group and emphasize related ideas. If you have more than one bulleted list in your report or presentation, the symbol you choose for the bullets should be consistent for all of your graphics. Use a minimum number of indent levels, providing more detail verbally. To avoid monotonous presentations, be careful not to overuse bulleted lists.

Pie Charts

A pie chart is the graphic that answers simple questions about proportions. Each slice represents an individual part of a particular group. "Cutting" (separating) one of the slices emphasizes an element that is part of the whole. For clarity, place labels next to the slices, not in a legend. Include percentages or values in the labels to add detail to the interpretation. Pie charts should contain eight slices at most. When you have more than eight data values, use a bar chart. Use multiple pie charts cautiously; bar charts are more effective in comparing proportions between groups.

2008 School Health Profiles Survey

Percentage of schools that require students to take the following number of required physical education courses.




Arrange your data from the largest element to the smallest, unless you want to emphasize a particular element, or there is a logical order to your categories or elements. Your most important element should start at the 3 o'clock position on the pie. The other elements should progress in importance in a counterclockwise direction, with each slice being a lighter color or shading. For the best color or pattern effects, work from dark to light. Fluctuating between dark and light makes it difficult to see pie shading differences.

Vertical Bar or Column Charts

Vertical bars are used to present trends in data such as changes over time or differences among groups. Use bar charts for a relatively small number of discrete data points or groups. Use a clustered bar chart to compare data in more than one category. However, keep the number of clusters small, and limit the number of bars in each cluster to three or fewer.


2008 School Health Profiles Survey

Percentage of schools that have a copy of the district wellness policy by school type



Horizontal Bar Chart

Horizontal bar charts are used to show comparisons between parts, groups, or categories. This type of chart will accommodate many values without visual clutter and can indicate exact quantities as well as proportions. Use the same color or fill pattern for all bars. To emphasize one bar, select a contrasting fill pattern or color. Your Profiles charts are in this format.



Line Charts

Line charts are used to show changes in data over time or to represent continuous measurements. Like bar charts, line charts answer questions about trends, and they can support an almost unlimited number of data points.

Titles and Labels

Graphics should have clear, concise titles and subtitles. Both axes of a graph should be labeled with the names of the variables, and the scales should be indicated. Titles should be centered at the top or bottom of the graphic. All information necessary to understand the graphic should be included.

Production

Graphics produced for paper copies and those created for computerized digital display require different design formats. It is important to consider the purpose and presentation medium when choosing among pattern, shading, and color options. Computer presentations benefit from use of color. Photocopying printed graphics (unless using a color copier) will obscure color or shading patterns.

Electronic Presentation

You can present your results in the PowerPoint presentation format provided without any modification, or you can tailor the report to meet more specific needs. The PowerPoint presentation allows you to add transitions between slides, text builds, and even animation and sound. Transitions are special visual effects that appear when moving from slide to slide. Text builds allow you to show main bullet points on a slide one at a time. These special effects should be used sparingly. You need to preview your presentation to be sure that everything looks good.

When doing an electronic presentation, you will need a computer, a CD or a flash drive with your "slides," and an LCD projector. In the future, LCD projectors connected to computers will probably become as commonplace as slide and overhead projectors. We are not there yet, however, so be prepared and bring your equipment. You may want to bring overheads or a printed version as a backup in case of equipment problems.

Internet Website

Presenting data on the Internet has become more commonplace. This format makes it readily available to many audiences. Having data available on the World Wide Web emphasizes the importance of the data and encourages an exchange of information to enhance analysis and presentation.

Preparing data for an Internet site is not difficult. Software packages are available to convert your paper presentation text and graphics into HTML (hypertext markup language) or PDF (portable document file), so it can be viewed on the Internet. Like your paper presentation, you will want to keep it simple and easy to read. Some formats will change in the HTML conversion, so take the time to adjust the format the way you like. Highlight important headings and keep basic facts bulleted. Converting to HTML will allow you to draw attention to links including appendices, tables, graphs, and pie charts, if you choose. By using hyperlinks, you can allow the user to move within your report or to jump to supplemental information available elsewhere on the Internet. Converting to PDF format will keep your document true to the original format.

Quality Assurance

Quality assurance is the time and effort spent by the graphics developer to ensure that the message conveyed by the graphic is true to the data it represents. Adequate quality assurance ensures that a graphic represents data in a manner that is easily viewed and understood by the observer and is not in any way misleading or incorrect. If the graphic presentation is incorrect, the viewer will have a false sense of the data and their implications.

Proofread your charts. If possible, enlist one or more of your co-workers who are familiar with your Profiles to help with the proofing. Also, check that percentages sum to 100%, when applicable, and that counts sum to the total. If percentages do not sum to 100 due to rounding, be sure to document that in a footnote.

Make sure the numbers on your chart match the numbers in the original data and that they are presented in the correct category. Within a presentation, scale changes should be avoided whenever possible so that between-chart comparisons can be made. For example, you may have two charts side by side showing response rates. One may use a scale of 0 to 100. The second chart may zoom in on a scale of 60 to 100. A person comparing these two charts will probably get a distorted view of the data. If you need to enlarge a selected portion of a scale, be sure it is clearly labeled as such. The vertical scale of bar and line charts should include zero.

Answer the following questions when proofreading your charts:

- Is all the text there? (Did the computer truncate text on long lines?)
- Is the spelling correct? (If your graphics package has one, use the built-in spell checker.)
- Is your message clear?
- Is the chart simple and easy to understand?
- Are the data accurate?
- Would color enhance the presentation of the data?

Whether you are compiling a written report or preparing visuals for a presentation, graphics can be used to add emphasis to your message. Graphics can help make sure your readers or audiences leave with the message you want to convey. Effective use of graphics may help you generate interest in your program, gain support for conducting Profiles, and enhance your report or presentation.

Checklist for Effective Graphics

Purpose

- Identify your audience(s).
- Specify your objectives.
- Ensure presentation methods match purpose and audience.

Planning

- Create rough drafts first.
- Plan on making several drafts of all graphs.
- Remember that producing graphics sometimes takes longer than expected, so plan time accordingly.

Appropriate Use

- Use graphics to highlight the intended material.
- Use the correct type of chart for your data.
- Be sure the chart demonstrates the comparisons you planned.

Clarity

- Avoid unnecessary shadowing, 3D effects, and coloring.
- Minimize the number of fonts.
- Use bold and italic versions of fonts for highlighting.
- Avoid red and green adjacent to each other.
- Use accurate and complete labels.

Simplicity

- Present the data without extraneous material.
- Avoid elaborate fill patterns.
- Avoid too many different patterns.
- Avoid overly decorative backgrounds.

Consistency

- Use a similar style across all graphics.
- Use comparable scales for accurate comparison.

Accuracy

- Check that data are correct.
- Check that spelling is correct.
- Double-check everything!

Washington
2008 School Health Profiles Report
PRINCIPAL QUESTIONNAIRE ITEM RATIONALE

Question 1: Are any of the following grades taught in this school?

Rationale: Based on the response to this question, the eligibility of the school is determined.

Question 2: Has your school ever used the School Health Index or other self-assessment tool to assess your school's policies, activities, and programs in the following areas?

Rationale: This question assesses whether the school has conducted an assessment or diagnosis as a critical first step in improving implementation of policies, programs, or environmental strategies to effect change or improvement in school health (1). Studies confirm that the School Health Index helps bring health issues to the school's attention, builds school commitment, identifies changes that do not require resources, encourages development of policy and action, raises awareness of federal policies, and helps schools set policies and standards that meet national health objectives (2,3).

Question 3: The Child Nutrition and WIC Reauthorization Act of 2004 requires school districts participating in federally subsidized child nutrition programs (e.g., National School Lunch Program, School Breakfast Program) to establish a local school wellness policy. Do you have a copy of your district's wellness policy?

Rationale: This question assesses school-level awareness of the district's wellness policy. According to the Child Nutrition and WIC Reauthorization Act of 2004, each school division that participates in the National School Lunch Program (NSLP) shall establish a local (school) wellness policy no later than the first day of the school year beginning after June 30, 2006, to cover all NSLP schools in the school division (4). It is important to measure school-level awareness of local wellness policies in order for state education and health agencies to provide technical assistance on the development, implementation, and evaluation of these policies, given the potential impact of school health policies on physical activity, diet, and the availability of foods and beverages (5,6).

Question 4: Currently, does someone at your school oversee or coordinate school health and safety programs and activities?

Rationale: This question assesses whether the school has identified a person responsible for coordinating a school's health program. It is critical to have one person appointed to oversee the school health program (7). This individual coordinates school health activities, leads a school health committee or team, and integrates community-based programs with school-based programs (8,9) Administration and management of school health programs requires devoted time, attention, training, and expertise (10,11).

Question 5: Is there one or more than one group (e.g., a school health council, committee, or team) at this school that offers guidance on the development of policies or coordinates activities on health topics?

Question 6: Are each of the following groups represented on any school health council, committee, or team?

Rationale: These questions assess whether the school has a health committee or team. The school health committee or team should represent a coalition of representatives from within and outside of the school community interested in improving the health of youth in schools (10,12). Participation on such committees or teams can empower others through increased awareness and knowledge of the school health program, increase the chance of ownership and commitment, activate channels of communication, and increase involvement in decision making (8,10,12-16). This includes parents and community members. Parent leaders help other parents understand and contribute ideas to issues and policies that affect the design and quality of school programs and opportunities for all children (15).

Question 7: Are any school staff required to receive professional development (such as workshops, conferences, continuing education, or any other kind of in-service) on HIV, STD, or pregnancy prevention issues and resources for the following groups?

Rationale: This question assesses professional development requirements for school staff. Professional development enhances the health educator's ability to teach adolescents the knowledge and skills needed to prevent HIV, other STDs,

and pregnancy. In the 2000 School Health Policies and Program Study, only 35% of health education teachers reported receiving staff development on HIV prevention during the two preceding years (17). Educators who have received professional development in health education report increases in the number of health lessons taught and their confidence in teaching (18). Professional development increases educators' confidence in teaching subject matter and provides opportunities for educators to learn about new developments in the field and innovative teaching techniques, and to exchange ideas with colleagues (7,19).

Question 8: Does this school have a student-led club that aims to create a safe, welcoming, and accepting school environment for all youth, regardless of sexual orientation or gender identity? These clubs sometimes are called gay/straight alliances.

Rationale: This question assesses whether the school has a gay/straight alliance or similar student-led club. Such clubs are critical to the well-being of students. Students in schools with a gay/straight alliance are less likely to feel unsafe at school, less likely to miss school, and more likely to feel like they belonged at their school than students in schools with no such clubs (20).

Question 9: Has your school adopted a policy that addresses each of the following issues for students or staff with HIV infection or AIDS?

Rationale: This question assesses important components of school policies in place to address students and staff infected with HIV or AIDS. Students and staff infected with HIV or AIDS need policies protecting their rights (21).

Question 10: Are all staff who teach health education topics at this school required to be certified, licensed, or endorsed by the state in health education?

Rationale: This question addresses the necessary qualifications of staff who teach health education. The National Commission for Health Education Certification, Inc. supports Certified Health Education Specialists (CHES). CHES teachers are more likely than non-CHES teachers to teach about HIV and sexually transmitted disease prevention (22). Studies in other disciplines reveal that nationally certified teachers, to a greater degree than non-certified teachers, possess pedagogical content knowledge that is more flexibly and innovatively employed in instruction; are more able to improvise and to alter instruction in response to contextual features of the classroom situation; understand at a deeper level the reasons for individual student success and failure on any given academic task; are more able to provide developmentally appropriate learning tasks that engage, challenge, and even intrigue students; are more able to anticipate and plan for difficulties students are likely to encounter with new concepts; are more easily able to improvise when things do not run smoothly; are more able to generate accurate hypotheses about the causes of student success and failure; and bring a distinct passion to their work (23). The leading national organizations supporting school health education, including the American School Health Association and the American Association for Health Education, recommend that those who teach health education have professional preparation and state certification in health education (24,25). The Joint Committee on National Health Education Standards recommends that local education agencies ensure that health education is taught by licensed/certified health education teachers with training in implementing the National Health Education Standards (26).

REQUIRED PHYSICAL EDUCATION

Question 11: Is physical education required for students in any of grades 6 through 12 in this school?

Question 12: Is a required physical education course taught in each of the following grades in this school?

Rationale: These questions measure the extent to which physical education is required for students in grades 6 through 12. Physical education provides students with the knowledge, attitudes, skills, behaviors, and confidence to adopt and maintain physically active lifestyles (27). The importance of physical education in promoting the health of young people is supported by *Healthy People 2010* Objectives 22-8, 22-9, and 22-10 (28).

Question 13: Can students be exempted from taking required physical education for one grading period or longer for any of the following reasons?

Rationale: This question examines whether or not students are allowed to be exempt from physical education based upon participation in non-physical activities or interscholastic sports. Exemptions from required physical education do not allow students to participate in comprehensive, standards-based physical education; this practice diminishes the

importance of physical education and its role in assisting students with establishing physically active lifestyles and developing various motor, movement, and behavioral skills unique to being physically educated (29).

PHYSICAL EDUCATION AND PHYSICAL ACTIVITY

Question 14: Are all staff who teach physical education at this school certified, licensed, or endorsed by the state in physical education?

Rationale: This question measures the necessary qualifications of staff who teach physical education (PE). The National Association for Sport and Physical Education (NASPE) recommends that those who teach PE have grade-specific preparation (30). CDC's *Guidelines for School and Community Programs to Promote Lifelong Physical Activity Among Young People* recommend that schools require the hiring of physical education specialists to teach physical education in kindergarten through grade 12 (31). Staff who have qualifications for teaching PE keep students active longer during PE classes and are more likely to teach to national, state, and/or local PE standards (32,33).

Question 15: During the past two years, did any physical education teachers or specialists at this school receive professional development (such as workshops, conferences, continuing education, or any other kind of in-service) on physical education?

Rationale: This question examines professional development for PE teachers. PE teachers who participate in staff development programs are more likely to use recommended teaching methods such as holding group discussions, implementing physical activity stations, videotaping student performances, testing students' knowledge related to PE, giving fitness tests, and explaining to students the meaning of fitness scores (30,34,35). Professional development for PE teachers provides skills to increase the quality of PE classes through student engagement in physical activity and the content of lessons taught (32,36,37).

Question 16: Are those who teach physical education at this school provided with the following materials?

Rationale: This question measures the type of information and support materials PE teachers are given in order to implement PE classes. According to NASPE, quality physical education is guided by and should include a written PE curriculum, goals, objectives, and expected outcomes, scope and sequence of instruction for PE, and plans for age-appropriate student assessment (34,38).

Question 17: Does this school offer opportunities for students to participate in intramural activities or physical activity clubs?

Rationale: This question measures the extent to which students are provided the opportunity to participate in physical activities and clubs outside of the regular school day. According to NASPE, intramural activities, physical activity clubs, and recreation clubs contribute to young people's physical and social development. Additionally, intramural activities or physical activity clubs offer students the opportunity to be involved in planning and implementing such programs and offer safe and structured opportunities to be physically active (39-43).

TOBACCO-USE PREVENTION POLICIES

Question 18: Has this school adopted a policy prohibiting tobacco use?

Question 19: Does the tobacco-use prevention policy specifically prohibit use of each type of tobacco for each of the following groups during any school-related activity?

Question 20: Does the tobacco-use prevention policy specifically prohibit tobacco use during each of the following times for each of the following groups?

Question 21: Does the tobacco-use prevention policy specifically prohibit tobacco use in each of the following locations for each of the following groups?

Question 22: Does your school have procedures to inform each of the following groups about the tobacco-use prevention policy that prohibits their use of tobacco?

Question 23: Does your school's tobacco-use prevention policy include guidelines on what actions the school should take when students are caught smoking cigarettes?

Question 24: At your school, who is responsible for enforcing your tobacco-use prevention policy?

Question 25: Which of the following help determine what actions the school takes when students are caught smoking cigarettes?

Question 26: When students are caught smoking cigarettes, how often is each of the following actions taken?

Question 27: Does your school post signs marking a tobacco-free school zone, that is, a specified distance from school grounds where tobacco use is not allowed?

Rationale: These questions measure the extent to which schools develop, implement, and enforce a policy that creates a totally tobacco-free environment within the school experience for both young people and adults, as outlined in the CDC *Guidelines for School Health Programs to Prevent Tobacco Use and Addiction* (44) to achieve the *Healthy People 2010* Objective 27-11 of creating smoke-free and tobacco-free schools. (28). The Pro-Children Act of 1994, reauthorized under the No Child Left Behind Act of 2001, prohibits smoking in facilities where federally funded educational, health, library, daycare or child development services are provided to children under the age of 18 (45). Because tobacco use is the most preventable contributor to mortality in the United States, it is important to restrict use or exposure to tobacco products at an early age (44). The existence and enforcement of a school policy creates a tobacco-free environment that models acceptable behavior and sends a clear message to students, teachers, staff, parents, and visitors that the use of tobacco is socially unacceptable (46). Environmental interventions aimed at reducing use of tobacco in homes, public places, and worksites lead to reduction of tobacco use (47). Likewise, tobacco-free school policies are associated with lower rates of student smoking (46,48-50). Prohibiting any use of any tobacco product at all times, whether or not school is in session, and regardless of whether students are present, protects students and staff from the harmful effects of secondhand smoke (a mixture of smoke from the burning end of tobacco products and the smoke exhaled by smokers). The 2006 U.S. Surgeon General's report, *The Harmful Effects of Involuntary Exposure to Tobacco Smoke*, outlines a large body of research findings which demonstrate that breathing secondhand smoke is harmful to health (51). Evidence shows that there is no safe level of secondhand smoke exposure, and even the most advanced ventilation systems cannot eliminate secondhand smoke or its harmful effects (51). A complete ban of indoor smoking *at all times* in a facility (such as a school building) is the only effective approach to controlling involuntary inhalation of secondhand smoke (51). Therefore, schools should eliminate smoking in school facilities by all persons at all times even when students or staff are not present. Additionally, preventing exposure to secondhand smoke at all school events and in school vehicles is also imperative to protect the health of all.

Question 28: During the past two years, has your school...Gathered and shared information with students and families about mass-media messages or community-based tobacco-use prevention efforts? Worked with local agencies or organizations to plan and implement events or programs intended to reduce tobacco use?

Rationale: These questions measure the extent to which the school coordinates their efforts with other tobacco-use prevention efforts in the community that target young people. School programs alone can be effective, but maintaining those effects present a challenge, especially with the many other influences encouraging tobacco use coming from outside of the child's school environment. The strongest evidence of success for school-based education programs has been shown with programs that are coordinated or delivered in conjunction with mass media and community tobacco control efforts, creating an environment of support for a tobacco-free lifestyle and delivering messages that are mutually reinforced. The Surgeon General has reported that 20-40% of tobacco use by youth can be prevented by educational strategies conducted in conjunction with community-and media-based activities (52). In 2004, the Task Force for the Community Guide to Preventive Services found "strong evidence for the use of school-based interventions when delivered in conjunction with mass media and community activities" (53). The Task Force recommendations were based on a subset of studies that showed evidence in reducing tobacco use among youth when multiple channels were used to support the in-school education program. Receiving consistent messages across community contexts and over time has been shown to enhance the maintenance of programs effects (54,55).

Question 29: Does your school provide tobacco cessation services for each of the following groups?

Question 30: Does your school have arrangements with any organizations or health care professionals not on school property to provide tobacco cessation services for each of the following groups?

Rationale: These questions measure the extent to which schools provide access to tobacco-use cessation services, as outlined in the CDC *Guidelines for School Health Programs to Prevent Tobacco Use and Addiction* (44) to achieve the *Healthy People 2010* Objectives 27-5 and 27-7 of increasing tobacco-use cessation attempts among adult and adolescent smokers (52). Nicotine addiction can occur at an early age for some adolescent tobacco users (56). People who begin using tobacco at an early age are more likely to develop higher levels of addiction in adulthood (56). Adolescent tobacco users suffer similar symptoms of withdrawal to those of adults when attempting to quit (57). Many young people want to quit but have tried and failed (58). Some are unaware of or do not have access to cessation services. Others

underestimate the power of addiction and do not feel that quitting would require professional assistance; therefore recruitment into formal programs can be difficult (59). School health providers as a routine part of care should assess the tobacco-use status of students, and if they identify a student's use of tobacco, they should provide self-help materials and refer them to a tobacco-use cessation program provided on site or in the community (60-62). Also, providing a brief clinical intervention has been shown to encourage cessation among both adults and adolescents (62).

NUTRITION-RELATED POLICIES AND PRACTICES

Question 31: When foods or beverages are offered at school celebrations, how often are fruits or non-fried vegetables offered?

Question 32: Can students purchase snack foods or beverages from one or more vending machines at the school or at a school store, canteen, or snack bar?

Question 33: Can students purchase each of the following snack foods or beverages from vending machines or at a school store, canteen, or snack bar?

Question 34: Does this school limit the package or serving size of any individual food and beverage items sold in vending machines or at the school store, canteen, or snack bar?

Rationale: These questions address the extent to which schools are making more nutritious foods available to students, limiting portion sizes, and not offering less nutritious foods and beverages. Many schools offer foods and beverages in after-school programs, school stores, snack bars, or canteens (63) and these foods sold in competition to school meals are often relatively low in nutrient density and relatively high in fat, added sugars and calories (64). Competitive foods are widely available in many elementary schools, in most middle schools, and in almost all secondary schools (63,65,66). Given that schools offer numerous and diverse opportunities for young people to learn and make consumption choices about healthful eating, schools should provide a consistent environment that is conducive to healthful eating behaviors (67). To help improve dietary behavior and reduce overweight among youths, schools should offer appealing and nutritious foods in school snack bars and vending machines and discourage sale of foods high in fat, sodium, and added sugars, and beverages and foods containing caffeine on school grounds (68-70). Because students' food choices are influenced by the total food environment, the simple availability of healthful foods such as fruits and vegetables may not be sufficient to prompt the choice of fruits and vegetables when other high fat or high sugar foods are easily accessible (71,72). However, offering a wider range of healthful foods can be an effective way to promote better food choices among high school students (73). Taken together, such findings suggest that restricting the availability of high-calorie, energy dense foods in schools while increasing the availability of healthful foods might be an effective strategy for promoting more healthful choices among students at school (67).

Question 35: During this school year, has your school done any of the following?

Rationale: This question addresses the variety of methods schools can use to promote healthy eating. Students' food choices are influenced by the total food environment. The simple availability of fruits and vegetables may not be sufficient to prompt the choice of these items when items high in fat and/or added sugar are also available (72). Even when fruit and vegetable items are available, they compete in the context of a vast array of other food items, mostly high in fat and sugar, that are competitively priced (71). Schools should employ effective or promising strategies in the school setting to promote healthy eating, such as pricing strategies (74,75), input from stakeholders (76), provision of nutrition information (67), taste tests, and using the cafeteria as a learning laboratory (77).

Question 36: At this school, are candy, meals from fast food restaurants, or soft drinks promoted through the distribution of products, such as t-shirts, hats, and book covers to students?

Question 37: Does this school prohibit advertisements for candy, fast food restaurants, or soft drinks in the following locations?

Rationale: These questions addresses prohibiting marketing of less nutritious foods to students while at school or at school-sponsored events. Nationwide, 23.3% of schools allow the promotion of candy, meals from fast food restaurants, or soft drinks through the distribution of coupons for free or reduced price, 14.3% allow the promotion of these products through sponsorship of school events, and 7.7% do so through publications such as a school newsletter or newspaper (63). Many contracts for soft drink or other vending products have provisions to increase the percentage of profits schools receive when sales volume increases, and this is a substantial incentive for schools to promote soft drink

consumption by adding vending machines, increasing the times they are available, and marketing the products to students (64,78). In some districts, these incentives have led schools to aggressively promote student purchases of soft

drinks (79). Research suggests that exposure to advertisements may have adverse impacts on children's eating habits (80). Food advertisements have been found to trigger food purchase by parents, have effects on children's product and brand preferences, and have an effect on consumption behavior (81). Further, younger children do not generally understand the difference between information and advertising, (82) such that children may interpret school-based advertising to mean that teachers or other adults endorse the use of the advertised product. Given that schools provide a captive audience of students, the IOM Report on Food Marketing to Children and Youth recommends that schools should promote healthful diets for children and youth in all aspects of the school environment (e.g., commercial sponsorships, meals and snacks, curriculum), and outlines the importance of prohibiting advertising of less nutritious foods (83).

HEALTH SERVICES

Question 38: Is there a full-time registered nurse who provides health services to students at your school?

Rationale: This question examines the degree to which schools are being adequately staffed by school nurses. Because a school nurse is an essential component of a healthy school, *Healthy People 2010* Objective 7-4 calls to increase the proportion of elementary, middle, and senior high schools with a nurse-to-student ratio of 1:750 (28). School nurses can link students and schools to physician and community resources.

Question 39: Which of the following sources of school health information does your school use to identify students diagnosed with chronic health conditions such as asthma?

Rationale: This question examines the appropriate use or adaptation of existing school health records to identify students with asthma. Asthma is a common chronic condition and one of the leading causes of school absenteeism. In 2003, students who had asthma episodes missed an estimated 12.8 million days of school (84). Asthma can be controlled with proper diagnosis, appropriate asthma care, and management activities. Schools play an important role in these asthma management activities, but individuals must first be identified in order to receive school-based support (85). Successful school-based asthma management programs identify students with asthma utilizing low-cost and/or no-cost adaptations of existing school health records and by avoiding mass screening and mass case detection as methods for routine identification of students with asthma (86-91).

Question 40: At your school, how many students with known asthma have an asthma action plan on file?

Rationale: This question addresses the need for clear, written guidance about the needs of individual students with asthma. Assessment of successful school-based asthma management programs suggest these plans play an important role in providing school staff, students, and families with an understanding of an individual student's asthma management needs at school, including how to respond in an emergency. Additionally, the use of an asthma action plan at school results in affected students experiencing significant improvement in several health-related outcomes, including a decrease in the frequency of asthma-related nighttime awakenings, number of days of restricted activity, and frequency of acute medical treatment (92,93). Schools should have asthma action plans on file for all students with known asthma. These plans help schools meet the needs of students with asthma during the school day and at school-related activities. Based upon current research, federal agencies and other national organizations have provided additional guidance and recommendations related to the collection and implementation of individualized plans. Plans should be developed by a primary care provider and be provided by parents; and include individualized emergency protocol, medications, environmental triggers and emergency contact information. School staff should actively solicit copies of asthma action plans from families and/or asthma care providers. When necessary, school nurses can construct asthma action plans based on information from the family and medication orders. A constructed plan should be sent to the asthma care provider for confirmation that it is appropriate (85,94-96).

Question 41: At your school, which of the following information is used to identify students with poorly controlled asthma?

Rationale: This question examines the type of information schools use to monitor and then assess the need for additional case management of students with known asthma. Assessment of successful school-based asthma management programs reveal that this type of tracking and case management can contribute to the medical management of students with asthma (87,92,97,98). This information can subsequently be used by schools to focus their asthma programs on students with poorly managed asthma as demonstrated by frequent school absences, school health office visits, emergency department visits, or hospitalizations (85,99).

Question 42: Does your school provide the following services for students with poorly controlled asthma?

Rationale: This question examines whether schools provide intensive case management for students with poorly controlled asthma. Schools should ensure that case management is provided by a trained professional for students with frequent school absences, school health office visits, emergency department visits, or hospitalizations due to asthma (85,92,98,100-103). Assessment of successful school-based asthma management programs reveal that monitoring and then providing case management can contribute to the medical management of students with asthma (87,98). Case management activities help students better manage their asthma, and have been shown to decrease hospitalizations, emergency department visits, and school absences among students with severe, persistent, or poorly controlled asthma (97,104).

Question 43: Does this school have a designated and secure storage location for medications, including quick-relief asthma medications?

Question 44: Is this location accessible at all times by the school nurse or her designee?

Question 46: Has your school adopted a policy stating that students are permitted to carry and self-administer asthma medications?

Question 47: Does your school have procedures to inform each of the following groups about your school's policy permitting students to carry and self-administer asthma medications?

Question 48: At your school, who is responsible for implementing your school's policy permitting students to carry and self-administer asthma medication?

Rationale: These questions address the need for schools to have policies and procedures to support students in receiving the asthma medications they may need at school. Many students with asthma require preventive or quick-relief medicine at school. Students with asthma have had serious episodes and have died at school when they did not have access to quick-relief medicine (105). Access to medications is critical and it must meet usual safety guidelines for medication storage (94,106). To ensure compliance with federal, state, and many local laws and guidelines, schools should ensure that students have immediate access to asthma medications, as prescribed by a physician and approved by parents (85). Several national guidance documents, along with evaluations of successful school based asthma programs, have provided additional information that addresses the process and methods for self-carry policies. Policies should include medication storage in a safe, controlled, and accessible location, and appropriate attention should be given to expiration dates and safe disposal (95,107-109).

Question 45: How often are school staff members required to receive training on recognizing and responding to severe asthma symptoms?

Rationale: This question examines professional development for school staff. Because asthma can be life-threatening, it is essential to assist those involved in monitoring and managing children with asthma at school to provide timely, appropriate care. Therefore, all school staff members should be provided with basic information about asthma so that they can support students' asthma management and appropriately respond to asthma emergencies (85,87,92,95,101,103,107).

FAMILY AND COMMUNITY INVOLVEMENT

Question 49: During the past 2 years, have students' families helped develop or implement policies and programs related to the following topics?

Question 50: During the past 2 years, have community members helped develop or implement policies and programs related to the following topics?

Rationale: These questions emphasize the importance of engaging family and community members in school health programs. Parent leaders help other parents understand and contribute ideas to issues and policies that affect the design and quality of school programs and opportunities for all children (16). School programs that engage parents and link with the community yield stronger positive results. Studies aimed at preventing childhood overweight, treating childhood overweight, or promoting physical activity and healthy eating have demonstrated more success when targeting the parent and child versus targeting the child alone (110,111). School-based tobacco prevention programs and community interventions involving parents and community organizations have a stronger impact over time when working in tandem rather than as separate, standalone interventions (55). Collaboration with parent groups, community organizations, and other agencies can help to build broad-based support for school health programs, especially when they address topics that can be emotionally charged, such as HIV, other STD, and pregnancy prevention (112). Without parental support of HIV, other STD, and pregnancy prevention education programs and policies, they cannot be sustained (112-114).

Collaborative asthma interventions require a team effort and involve the whole school community: school administrators, faculty, and staff, as well as students, parents, and local community organizations (95,103).

REFERENCES

1. Goodman R, Steckler A, Kegler MC. Mobilizing organizations for health enhancement. In: Glantz K, Lewis FM, Rimer B. eds. *Health Behavior and Health Education*. San Francisco, CA: Jossey Bass Publishers, 1997, pp. 287-312.
2. Pearlman DN, Dowling E, Bayuk C, Cullinen K, Thacher AK. From concept to practice: using the School Health Index to create healthy school environments in Rhode Island elementary schools. *Preventing Chronic Disease* [serial online] 2005; Nov.
3. Staten LK, Teufel-Shone NI, Steinfelt VE, et al. The School Health Index as an impetus for change. *Preventing Chronic Disease* [serial online] 2005; Jan.
4. US 108th Congress, Child Nutrition and WIC Reauthorization Act of 2005. S 2507. Available at: <http://www.govtrack.us/congress/bill.xpd?bill=s108-2507>
5. French SA, Story M, Fulkerson JA, Gerlach AF. Food environment in secondary schools: a la carte, vending machines, and food policies and practices. *American Journal of Public Health* 2003; 93:1161-1167.
6. Davee AM, Blum JE, Devore RL, et al. The vending and a la carte policy intervention in Maine public high schools. *Preventing Chronic Disease* [serial online] 2005; 2 Spec.
7. Institute of Medicine. *Schools and Health: Our Nation's Investment*. Washington, DC: National Academy Press, 1997.
8. Fetro JV. Implementing Coordinated School Health Programs in Local Schools. In: Marx E, Wooley S, eds. *Health Is Academic: A Guide to Coordinated School Health Programs*. New York: Teachers College Press, 1998.
9. American Cancer Society. *School Health Program Elements of Excellence: Helping Children to Grow Up Healthy and Able to Learn*. Atlanta, GA: American Cancer Society; 2000.
10. National Association of State Boards of Education. *Fit, Healthy, and Ready to Learn: A School Health Policy Guide*. National Association of State Boards of Education. Washington DC: NASBE, 2000.
11. American Cancer Society. *Improving School Health: A Guide to the Role of School Health Coordinator*. Atlanta, GA: American Cancer Society, 1999.
12. Shirer K. *Promoting health Youth, Schools and Communities: A Guide to Community-School Health Councils*. Atlanta, GA: American Cancer Society, 2003.
13. Green, LW, Kreuter MW. *Health Promotion and Planning: An Education and Environmental Approach*. California: Mayfield Publishing Company; 1991, pp. 271-274.
14. Birch, DA. Involving families in school health education: an essential partnership. In: Cortese P, Middleton K, eds. *The Comprehensive School Health Challenge: Promoting Health Through Education*. Vol 1. Santa Cruz, CA: ETR Associates, 1994.
15. Redding, S. et.al. *The Effects of Comprehensive Parent Engagement on Student Learning Outcomes*. Presentation at the Annual Convention of American Educational Research Association, San Diego, 2004.
16. Epstein LS. *School, Family, and Community Partnerships: Preparing Educators and Improving Schools*. Boulder, CO: Westview Press, 2001.
17. Kann L, Brener N, Allensworth D. Health Education: Results from the School Health Policies and Programs Study 2000. *Journal of School Health* 2001; 71(7):266-278.
18. Hausman A, Ruzek S.. Implementation of comprehensive school health education in elementary schools: focus on teacher concerns. *Journal of School Health* 1995; 65(3):81-86.
19. National Association of State Boards of Education (NASBE) & National School Boards Association (NSBA). *HIV Prevention in Schools: A Tool Kit for Education Leaders*. Alexandria, VA, 2002. Available at: http://www.nasbe.org/healthy_schools/publications/HIV_Tool_Kit.pdf
20. Kosciw JG, Diaz EM. *The 2005 National School Climate Survey: The experiences of lesbian, gay, bisexual, and transgender youth in our nation's schools*. New York: Gay, Lesbian, and Straight Education Network, 2006.
21. National Association of State Boards of Education. *Someone at School Has AIDS: A Comprehensive Guide to Education Policies Concerning HIV Infection*. Alexandria, VA: National Association of State Boards of Education, 1996.
22. Jones SE, Brener ND, McManus T. The relationship between staff development and health instruction in schools in the United States. *American Journal of Health Education* 2004; 35:2-
23. Bond L, Smith T, Baker WK, Hattie JA. The certification system of the National Board for Professional Teaching Standards: A construct and consequential validity study. University of North Carolina at Greensboro, Center for Educational Research and Evaluation, 2000. Available at: http://www.nbpts.org/resources/research/browse_studies?ID=168
24. American Association for Health Education. A Position Statement: Certification of Health Education Teachers, 2005. Available at: AAHE@aahperd.org.
25. American School Health Association. Resolution: Certification, 2002. Available at: <http://www.ashaweb.org/pdfs/resolutions/Certification.pdf>.
26. The Joint Committee on National Health Education Standards. *National Health Education Standards: Achieving Excellence (2nd Edition)*. Atlanta: American Cancer Society, 2007.
27. Burgeson C, Wechsler H, Brener N, Young J, Spain C. Physical education and activity: results from the School Health Policies and Programs Study 2000. *Journal of School Health* 2001; 71:279-293.

28. U.S. Department of Health and Human Services. *Healthy People 2010*. 2nd ed. With *Understanding and Improving Health and Objectives for Improving Health*, 2 vols. Washington, DC: U.S. Department of Health and Human Services, 2000.
29. National Association for Sport and Physical Education. *Opposing substitution and waiver/exemption for required physical education*. Reston, VA: National Association for Sport and Physical Education, 2006. Available at www.aahperd.org/naspe.
30. National Association for Sport and Physical Education. *National standards for beginning physical education teachers*. Reston, VA: National Association for Sport and Physical Education, 2001.
31. CDC. Guidelines for school and community programs to promote lifelong physical activity among young people. *MMWR* 1997;46(No. RR-6).
32. McKenzie TL, Feldman, H, Woods SE, Romero KA, Dahlstrom V, Stone EJ, et al. Children's activity levels and lesson context during third-grade physical education. *Research Quarterly for Exercise and Sport* 1996; 66(3):184-193.
33. Sallis JF, McKenzie TL, Alcaraz JE, Kolody B, Faucette N, Hovell MF. The effects of a 2-year physical education program (SPARK) on physical activity and fitness in elementary school students. *Sports, Play, and Active Recreation for Kids. American Journal of Public Health* 1997; 87(8):1328-1334.
34. National Association for Sport and Physical Education. *Moving into the future: National standards for physical education*. Reston, VA: National Association for Sport and Physical Education, 2004.
35. Davis K, Burgeson CR, Brener ND, McManus T, Wechsler H. The relationship between qualified personnel and self-reported implementation of recommended physical education practices and programs in U.S. schools. *Research Quarterly for Exercise and Sport* 2005; 76(2):202-211.
36. Kelder S, Mitchell PD, McKenzie TL, Derby C, Strikmiller PK, Luepker RV, et al. Longterm implementation of the CATCH physical education program. *Health Education and Behavior* 2003; 30(4):463-475.
37. McKenzie TL, Marshall SJ, Sallis JF, Conway TL. Student activity levels, lesson context, and teacher behavior during middle school physical education. *Research Quarterly for Exercise and Sport* 2000; 71(3):249-259.
38. National Association for Sport and Physical Education. *What constitutes a quality physical education program?* Reston, VA: National Association for Sport and Physical Education, 2003. Available at www.aahperd.org/naspe.
39. National Association for Sport and Physical Education. *Guidelines for after-school physical activity and intramural programs*. Reston, VA: National Association for Sport and Physical Education, 2002. Available at www.aahperd.org/naspe.
40. Hellison D. Physical activity programs for underserved youth. *Journal of Science & Medicine in Sport* 2000; 3(3):238-42.
41. Kelder S, Hoelscher DM, Barroso CS, Walker JL, Cribb P, Hu S. The CATCH Kids Club: a pilot after-school study for improving elementary students' nutrition and physical activity. *Public Health Nutrition* 2005; 8(2):133-40.
42. Pate RR, Saunders RP, Ward DS, Felton G, Trost SG, Dowda M. Evaluation of a community-based intervention to promote physical activity in youth: lessons from Active Winners. *American Journal of Health Promotion* 2003; 17(3):171-82.
43. Trevino RP, Yin Z, Hernandez A, Hale DE, Garcia OA, Mobley C. Impact of the Bienestar school-based diabetes mellitus prevention program on fasting capillary glucose levels: a randomized controlled trial. *Archives of Pediatrics & Adolescent Medicine* 2004; 158(9):911-7.
44. CDC. Guidelines for school health programs to prevent tobacco use and addiction. *MMWR* 1994; 43(No. RR-2).
45. Pro-Children Act of 1994, 20 U.S.C. §6081-6084 (1994).
46. Brownson RC, Koffman DM, Novotny TE, Hughes RG, Eriksen MP. Environmental and policy interventions to control tobacco use and prevent cardiovascular disease. *Health Education Quarterly* 1995; 22(4):478-98.
47. Levy DT. The effects of tobacco control policies on smoking rates: a tobacco control scorecard. *Journal of Public Health Management and Practice* 2004; 10(4):338-53.
48. Wakefield MA, Chaloupka FJ, Kaufman NJ, Orleans CT, Barker DC, Ruel EE. Effect of restrictions on smoking at home, at school, and in public places in teenage smoking: cross sectional study. *British Medical Journal* 2000; 321:310-311.
49. Charlton A, While D. Smoking prevalence among 16-19 year olds related to staff and student smoking policies in sixth forms and further education. *Health Education Journal* 1994; 53:191-215.
50. Pentz MA, Brannon BR, Carlin VL, Barrett EJ, MacKinnon DP, Flay BR. The power of policy: the relationship of smoking policy to adolescent smoking. *American Journal of Public Health* 1989; 79:857-62.
51. U.S. Department of Health and Human Services. *The Health Consequences of Involuntary Exposure to Tobacco Smoke: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, Coordinating Center for Health Promotion, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2006.
52. U.S. Department of Health and Human Services. *Reducing Tobacco Use: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.
53. Guide to Community Preventive Services. Task Force Meeting. February 25, 2004. Meeting minutes available at www.thecommunityguide.org.
54. Sussman S. School-based tobacco use prevention and cessation: where are we going? *American Journal of Health Behavior* 2001;25(3):191-9.
55. Lantz PM, Jacobson PD, Warner KE, Wasserman J, Pollack HA, Berson J, Ahlstrom A. Investing in youth tobacco control: a review of smoking prevention and control strategies. *Tobacco Control* 2000;9:47-
56. U.S. Department of Health and Human Services. *Preventing Tobacco Use Among Young People: A Report of the Surgeon General*. Atlanta, Georgia: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health; 1994.

57. Centers for Disease Control and Prevention. Reasons for Tobacco Use and Symptoms of Nicotine Withdrawal Among Adolescent and Young Adult Tobacco Users--United States, 1993. *Morbidity and Mortality Weekly Report* 1994, Vol. 43, pp. 746-750.
58. Eaton DK, Kann L, Kinchen S, et al. Youth risk behavior surveillance—United States, 2005. *Morbidity and Mortality Weekly Report* 2006; 55(No. SS-5).
59. Milton MH, Maule CO, Yee SL, Backinger C, Malarcher AM, Husten CG. Youth Tobacco Cessation: A Guide for Making Informed Decisions. Atlanta: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention; 2004.
60. Allensworth DD. Guidelines for Adolescent Preventive Services: a role for the school nurse. *Journal of School Health* 1996; 66(8):281-285.
61. Donovan KA. Smoking cessation programs for adolescents. *Journal of School Nursing* 2000; 16(4):36-43.
62. Fiore MC, Bailey WC, Cohen SJ, et. al. *Treating Tobacco Use and Dependence*. Quick Reference Guide for Clinicians. Rockville, MD: U.S. Department of Health and Human Services. Public Health Service, 2000.
63. Wechsler H, Brener ND, Kuester S, Miller C. Food service and foods and beverages available at school: results from the School Health Policies and Programs Study 2000. *Journal of School Health* 2001; 71(7):313-24
64. US Department of Agriculture. Foods sold in competition with USDA school meal programs: a report to congress. Food and Nutrition Service, 2001. Available at: http://www.fns.usda.gov/cnd/Lunch/CompetitiveFoods/report_congress.htm
65. Kann L, Grunbaum J, McKenna ML, Wechsler H, Galuska DA. Competitive foods and beverages available for purchase in secondary schools--selected sites, United States, 2004. *Morbidity and Mortality Weekly Report* 2005; 54(37):917-921.
66. US Government Accountability Office. School meal programs: Competitive foods are widely available and generate substantial revenues. Report to Congressional Requesters GAO-05-563, 2005. Available at: <http://www.gao.gov/new.items/d05563.pdf>.
67. Food and Nutrition Board, Institute of Medicine, Committee on Prevention of Obesity of Children and Youth--Schools. In: JP Koplan, CT Liverman, VI Kraak, (Eds.) *Preventing Childhood Obesity: Health in the Balance*. Washington, DC: National Academy Press, 2005, pp. 237–284.
68. Wechsler H, McKenna ML, Lee SM, Dietz WH. The role of schools in preventing childhood obesity. *The State Education Standard*, December 2004. Available at http://www.nasbe.org/standard/17_dec2004/wechsler2.pdf.
69. Pilant VB, American Dietetic Association. Position of the American Dietetic Association: Local Support for Nutrition Integrity in Schools. *Journal of the American Dietetic Association* 2006;106(1):122-33.
70. Institute of Medicine. *Nutrition Standards for Foods in Schools: Leading the Way Toward Healthier Youth*. Washington DC, Institute of Medicine of the National Academies, 2007.
71. Cullen KW, Eagan J, Baranowski T, Owens E, deMoor C. Effect of a la carte and snack bar foods at school on children's lunchtime intake of fruits and vegetables. *Journal of the American Dietetic Association* 2000; 100:1482–1486.
72. Kubik MY, Lytle LA, Hannan PJ, Perry CL, Story M. The association of the school food environment with dietary behaviors of young adolescents. *American Journal of Public Health* 2003; 93:1168–1173.
73. French SA, Story M, Fulkerson JA, Hannan P. An environmental intervention to promote lower fat food choices in secondary schools. Outcomes of the TACOS study. *American Journal of Public Health* 2004; 94(9):1507-1512.
74. French SA, Story M, Jeffery RW, Snyder P, Eisenberg M, Sidebottom A. Pricing strategy to promote fruit and vegetable purchase in high school cafeterias. *Journal of the American Dietetic Association* 1997; 97:1008–1010.
75. French SA, Jeffery RW, Story M, Breitlow KK, Baxter JS, Hannan PJ et al., Pricing and promotion effects on lowfat vending snack purchases: the CHIPS study. *American Journal of Public Health* 2001; 91:112–117.
76. Food and Nutrition Service US Department of Agriculture, Centers for Disease Control and Prevention, US Department of Health and Human Services, and US Department of Education. *Making It Happen: School Nutrition Success Stories*. Alexandria, VA: US Department of Agriculture, 2005.
77. American Dietetic Association, Position of the American Dietetic Association, Society for Nutrition Education, and American School Food Service Association. Nutrition services: an essential component of comprehensive school health programs, *Journal of the American Dietetic Association* 2003; 103: 505–514.
78. Square Meals: Nourishing Children's Bodies and Minds. School Vending Contract Survey. Available at: http://www.squaremeals.org/fn/render/channel/items/0,1249,2348_2515_0_0,00.html
79. Nestle M. Soft drink "pouring rights:" marketing empty calories to children. *Public Health Reports* 2000; 115:308-319.
80. Horgen KB, Choate M, Brownell KK. Television and food advertising: targeting children in a toxic environment. In: Sinder DG, Singer JL, eds. *Handbook of Children and the Media*. Thousand Oaks CA Sage Publications, 2001, pp. 447-461.
81. Hastings G, Stead M, McDermott L, Forsyth A, MacKintosh A, Rayner M, Godfrey C, Caraher M, Angus K. 2003. Review of Research on the Effects of Food Promotion to Children. Glasgow, UK. Center for Social Marketing, University of Strathclyde. Available at: <http://www.foodstandards.gov.uk/multimedia/pdfs/foodpromotiontochildren1.pdf>
82. Wilcox BL, Kunkel D, Cantor J, Dowrick P, Linn S, Palmer E. Report of the APA Task Force on Advertising and Children. Washington DC: American Psychological Association; 2004.
83. Committee on Food Marketing and the Diets of Children and Youth. *Food marketing to children and youth: threat or opportunity?* Washington, DC: Institute of Medicine of The National Academies Press, 2006.
84. Akinbami LJ. The state of childhood asthma, United States, 1980–2005. *Advance Data from Vital and Health Statistics*, no. 381, Hyattsville, MD: National Center for Health Statistics, 2006.
85. CDC. *Strategies for Addressing Asthma Within a Coordinated School Health Program, With Updated Resources*. Atlanta, Georgia: Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, 2006.
86. Boss LP, Wheeler LSM, Williams PV, et al. Population-based screening or case detection for asthma: are we ready? *Journal of Asthma* 2003; 40(4):335-342.

87. Splett PL, Erickson CD, Belseth SB, Jensen C. Evaluation and sustainability of the healthy learners asthma initiative. *Journal of School Health* 2006; 76(6):276-282.
88. Yawn BP. Asthma screening, case identification and treatment in school-based programs. *Current Opinion in Pulmonary Medicine* 2006; 12(1):23-27.
89. Yawn BP, Wollan P, Scanlon P, Kurland M. Are we ready for universal school-based asthma screening? An outcomes evaluation. *Archives of Pediatrics and Adolescent Medicine* 2002; 156(12):1256-1262.
90. Yawn BP, Wollan P, Scanlon P, Kurland M. Outcome results of a school-based screening program for under-treated asthma. *Annals of Allergy* 2003; 90:508-515.
91. Wheeler LS, Boss LP, Williams PV. School-based approaches to identifying students with asthma. *Journal of School Health* 2004; 74:378-380.
92. Erickson CD, Splett PL, Mullett SS, Jensen C, Belseth SB. The healthy learner model for student chronic condition management-Part II: The Asthma Initiative." *Journal of School Nursing* 2006; 22(6):319-329.
93. Gillies J, Barry D, Crane J, et al. A community trial of a written self management plan for children with asthma. *New Zealand Medical Journal* 1996; 109(1015):30-33.
94. National Association of School Nurses (NASN). Position Statement: Medication Administration in the School Setting. Revised June 2003. Available at: <http://www.nasn.org/Default.aspx?tabid=230>.
95. National Asthma Education and Prevention Program. Students with Chronic Illnesses: Guidance for Families, Schools and Students. National Heart, Lung, and Blood Institute, 2002. Available at: <http://www.nhlbi.nih.gov/health/public/lung/asthma/guidfam.htm>
96. National Asthma Education and Prevention Program. Managing Asthma: A Guide for Schools. Bethesda, Maryland: National Asthma Education and Prevention Program, National Heart, Lung, and Blood Institute, 2003. Available at: http://rover.nhlbi.nih.gov/health/prof/lung/asthma/asth_sch.htm
97. Levy M., Heffner B, Stewart T, Beeman G. The efficacy of asthma case management in an urban school district in reducing school absences and hospitalizations for asthma. *Journal of School Health* 2006; 76(6):320-324.
98. Taras H, Wright S, Brennen J, Campana J, Lofgren R. Impact of school nurse case management on students with asthma. *Journal of School Health* 2004; 74(6):213-219.
99. Barbot O, Platt R, Marchese C. Using preprinted rescue medication order forms and health information technology to monitor and improve the quality of care for students with asthma in New York City Public Schools. *Journal of School Health* 2006; 76(6):329-332.
100. National Association of School Nurses (NASN). School Nursing Management of Students with Chronic Health Conditions, 2006. Available at: <http://www.nasn.org/Default.aspx?tabid=351>
101. Taras H, Duncan P, Luckenbill D, Robinson J, Wheeler L, Wooley S, eds. *Health, Mental Health, and Safety Guidelines for Schools*. Elk Grove Village, Ill: American Academy of Pediatrics, 2004.
102. Tinkelman D, Schwartz A. School-based asthma disease management. *Journal of Asthma* 2004; 41(4):455-62.
103. Wheeler LS, Merkle SL, Gerald LB, Taggart VS. Managing asthma in schools: lessons learned and recommendations. *Journal of School Health* 2006; 76(6):340-344.
104. Evans R, Gergen PJ, Mitchell H, et al. A randomized clinical trial to reduce asthma morbidity among inner-city children: results of the National Cooperative Inner-City Asthma Study. *Journal of Pediatrics* 1999; 135(3):332-338.
105. Greiling AK, Boss LP, Wheeler LS. A preliminary investigation of asthma mortality in schools. *Journal of School Health* 2005; 75(8):286-290.
106. American Academy of Pediatrics, Committee on School Health. Guidelines for the Administration of Medication in School. *Pediatrics* 2003; 112(3):697-699.
107. National Asthma Education and Prevention Program. Resolution on Asthma Management at School. Updated November 2005. Available at: <http://www.nhlbi.nih.gov/health/public/lung/asthma/resolut.htm>.
108. National Asthma Education and Prevention Program. When Should Students with Asthma or Allergies Carry and Self-Administer Emergency Medications at School? Guidance for Health Care Providers Who Prescribe Emergency Medications. National Heart, Lung, and Blood Institute, 2005. Available at: http://www.nhlbi.nih.gov/health/prof/lung/asthma/emmer_medi.htm
109. National Association of School Nurses (NASN). The Use of Asthma Rescue Inhalers in the School Setting, Revised 2005. Available at: <http://www.nasn.org/Default.aspx?tabid=202>, 110. Golan M, Crow S. Targeting parents exclusively in the treatment of childhood obesity: long-term results. *Obesity Research* 2004; 12:357-361.
111. Epstein LH, Voloski A, Wing RR, McCurley J. Ten-year outcomes of behavioral family based treatment for childhood obesity. *Health Psychology* 1994; 13:373-83.
112. Council of Chief State School Officers. Joint Work Group. Essential tips for successful collaboration. Washington, DC: Author, 2004.
113. Council of Chief State School Officers. What Education Leaders Should Know About Forming Partnerships to Prevent Sexual-Risk Behaviors in School-Aged Youth. Washington, DC: Author, 2005.
114. Kirby D, Laris BA, Roller L. Sex and HIV education programs for youth: Their impact and important characteristics. Washington DC: Family Health International, 2006. Available at: www.etr.org/recapp/programs/SexHIVedProgs.pdf.

Washington
2008 School Health Profiles Report
LEAD HEALTH TEACHER QUESTIONNAIRE ITEM RATIONAL

REQUIRED HEALTH EDUCATION

Question 1: Is health education required for students in any of grades 6 through 12 in this school?

Question 2: How many required health education courses do students take in grades 6 through 12 in this school?

Question 3: Is a required health education course taught in each of the following grades in this school?

Rationale: These questions measure the extent to which health education courses are required for students in grades 6 through 12. School health education could be one of the most effective means to reduce and prevent some of the most serious health problems in the U.S., including cardiovascular disease, cancer, motor-vehicle crashes, homicide, and suicide (1). The Institute of Medicine (IOM) has recommended that U.S. schools require one-semester health education course at the secondary school level (1); however, the benefits of a health education curriculum increase when students receive at least three consecutive years of quality health curriculum (2).

Question 4: If students fail a required health education course, are they required to repeat it?

Rationale: This question measures the importance of a required health education course for students in grades 6 through 12.

Question 5: Are those who teach health education at this school provided with the following materials?

Rationale: This question addresses the types of information and support materials health education teachers are given in order to implement health education classes. According to the Joint Committee on National Health Education Standards, quality health education is guided by access and equity principles that call for clear curriculum direction, including goals, objectives, and expected outcomes; a written curriculum; clear scope and sequence of instruction for health education content; and plans for age-appropriate student assessment (3).

Question 6: Does your health education curriculum address each of the following?

Rationale: This question addresses the extent to which schools have a health education curriculum that is based on, or is consistent with, current national health education standards (3).

Question 7: During this school year, have teachers in this school tried to increase student knowledge on each of the following topics in a required course in any of grades 6 through 12?

Rationale: This question addresses the extent to which traditional health content areas and the prevention of health risk behaviors are taught in required courses in grades 6 through 12.

Question 8: During this school year, did teachers in this school teach each of the following tobacco-use prevention topics in a required course for students in any of grades 6 through 12?

Rationale: This question measures the tobacco-use prevention curricula content, and relates to the *Healthy People 2010* Objective 7-2 of providing school health education to prevent health problems among middle, junior high, and high school students including those from tobacco use (4). Since most smoking is initiated by persons less than 18 years old, programs that prevent onset of smoking during the school years are crucial (5). School-based tobacco prevention programs that address multiple psychosocial factors related to tobacco use among youth and that teach the skills necessary to resist those influences have demonstrated consistent and significant reductions or delays in adolescent smoking (5-12). Social influence programming has reduced smoking onset by as much as 50%, with effects lasting up to 6 years, and with effects including reduction of the use of other tobacco products as well (7). In addition, this question measures the extent to which schools are complying with the components of the National Health Education Standards, which provide a framework for decisions about the lessons, strategies, activities, and types of assessment to include in a health education curriculum (3).

Question 9: During this school year, did teachers in this school teach each of the following HIV, STD, or pregnancy prevention topics in a required course for students in any of grades 6, 7, or 8?

Question 10: During this school year, did teachers in this school teach each of the following HIV, STD, or pregnancy prevention topics in a required course for students in any of grades 9, 10, 11, or 12?

Rationale: These questions measure the HIV prevention curricula content. HIV and sex education programs can increase knowledge about how to avoid HIV and STD infection and unintended pregnancy. Efforts to increase knowledge alone about modes of transmission and strategies for prevention do not directly lead to behavior change (13). In order to reduce HIV, other STDs, and unintended pregnancy, programs must address perceptions of risk, intentions, communication, and skills, in addition to HIV and STD knowledge (14). Adolescents have different HIV prevention needs than do adults. However, because of the variability among youth with respect to cognitive and social maturity and sexual experience, interventions must be tailored to meet the unique needs of younger versus older youth or sexually naive versus experienced teens. HIV prevention interventions also have to be matched to the cognitive level of adolescents and should be designed to improve behavioral skills for risk reduction, decision making, planning, and problem solving (15). Part of a program's effectiveness involves its organization and presentation of activities and materials in an age appropriate and logical sequence. A typical logical sequence includes basic information about HIV, other STDs, and pregnancy, including susceptibility and severity of HIV, other STDs, and pregnancy; discussion of behaviors to reduce vulnerability; knowledge, values, attitudes and barriers related to these behaviors; and skills needed to perform these behaviors (16). Messages in these effective programs are appropriate to the age, sexual experience, gender and culture of the youth (14). In addition, the appropriateness for age and HIV prevention curricula content we realigned with the National Health Education Standards (NHES), which provide framework for decisions about the lessons, strategies, activities, and types of assessment to include in a health education curriculum (3). The NHES, other state-level standards, and research findings from the literature on effective programs in HIV, other STD, or pregnancy prevention were used to determine the activities and materials for age appropriateness for grades 6-8 and 9-12 related to HIV prevention curricula content (3,14-16).

Question 11: During this school year, did teachers in this school teach each of the following nutrition and dietary behavior topics in a required course for students in any of grades 6 through 12?

Rationale: This question measures the curricula content related to nutrition and dietary behavior. Comprehensive, sequential nutrition education using the classroom and the lunchroom can reinforce healthful eating behaviors (17,18). Nutrition education should be part of a comprehensive school health education curriculum and include concepts to promote healthy eating (19,20). This list of 15 nutrition topics is based on CDC guidelines (21) and the *School Health Index* (22).

Question 12: During this school year, did teachers in this school teach each of the following physical activity topics in a required course for students in any of grades 6 through 12?

Rationale: This question measures the extent to which physical activity concepts are taught in a required health education course. Health education that includes physical activity concepts increases the likelihood of students increasing their participation in physical activity (23), reinforces what has been taught in physical education (24,25), and assists students in achieving the national health education standards (3).

HIV PREVENTION

Question 13: During this school year, did your school provide any HIV, STD, or pregnancy prevention programs for ethnic/racial minority youth at high risk (e.g. black, Hispanic, or American Indian youth), including after-school or supplemental programs, that did each of the

following?

Rationale: This question measures whether a school addresses HIV, other STD, and pregnancy prevention through targeted efforts reaching those identified as most at-risk. Risk for HIV infection is especially notable for youth of minority races and ethnicities. African-Americans are the largest group of young people affected by HIV, accounting for 56% of all HIV infections ever reported among those aged 13–24 by 2001 (26). And, although only 15% of teenagers (ages 13-19) in the United States are African-Americans, they accounted for 73% of new AIDS cases reported among teens in 2004 (27). In 2002, HIV/AIDS was the number one cause of death for African-American women aged 25-34 years and the number two cause of death for all African-Americans aged 35-44 (28). Of the more than half a million people with AIDS who have died in the U.S., 38% were African American (29). In addition, the incidence of AIDS for adult and adolescent Latinos in 2001 was more than 3 times higher than that among their non-Hispanic white counterparts (29,30). Data from CDC's 2005 National Youth Risk Behavior Survey (YRBS) show that, compared with white students and Hispanic/Latino students, black students have the highest rates of several sexual risk behaviors: 67.6% of black students had ever had sexual

intercourse, compared with 43.0% percent of white students and 51.0% of Hispanic/Latino students; 47.4% of black students were currently sexually active (i.e., had sexual intercourse with 1 or more persons during the 3 months preceding the survey), compared with 32.0% of white students and 35.0% of Hispanic/Latino students; 16.5% of black students had had sexual intercourse before age 13 years, compared with 4.0% of white students and 7.3% of Hispanic/Latino students; and 28.2% of black students had sexual intercourse with 4 or more persons during their life, compared with 11.4% of white students and 15.9% of Hispanic/Latino students (31). In addition to effective curricula, access to valid information and products, as well as access or referral to health, social, and psychological services to prevent HIV, thursts, and pregnancy are especially important in ethnic/racial minority communities where the higher prevalence of HIV, other STDs, and pregnancy reflects both risky adolescent sexual behaviors and system barriers to quality prevention services (32). Factors which may influence adolescents' access to care include health insurance, cost, convenience, confidentiality, and demographic factors such as age, gender, and ethnicity (33-35).

COLLABORATION

Question 14: During this school year, have any health education staff worked with each of the following groups on health education activities?

Rationale: This question measure the extent to which health education staff work cooperatively with other components of the school health program (school health services, school mental health or social services, food service, and physical education staff) and families. An integrated school and community approach is an effective strategy to promote adolescent health and well being (36).

Question 15: During this school year, did your school provide parents and families with health information designed to increase parent and family knowledge of the following topics?

Rationale: This question measures whether schools are providing health information to students' families. School programs that engage parents and link with the community yield stronger positive results. Studies aimed at preventing childhood overweight, treating childhood overweight, and promoting physical activity and healthy eating have demonstrated more success when targeting the parent and child versus targeting the child alone (37,38). School-based tobacco prevention programs and community interventions involving parents and community organizations have a stronger impact over time when working in tandem rather than as separate, stand-alone interventions (10). Assessments of successful school-based asthma management programs indicate that with increased knowledge, parents can assist their children in better managing their asthma (39-41). Parents also are teenagers' primary sex educators, able to capitalize on teachable moments when youth may be more open to learning new information (42). Parents can continue prevention messages delivered in school, thereby enhancing the likelihood of sustained behavioral changes (43). Increased communication affects both parenting and health practices of parents. Communicating information on healthy lifestyles aims to reinforce the child's coursework at school, facilitate communication with parents about school activities, and increase parent knowledge of healthy living (44,45).

PROFESSIONAL DEVELOPMENT

Question 16: During the past two years, did you receive professional development (such as workshops, conferences, continuing education, or any other kind of in-service) on each of the following topics? (HIV)

Rationale: This question measures the extent to which professional development has been received by the lead health teacher responsible for teaching about HIV/AIDS. As new information and research on prevention is available, those responsible for teaching about HIV/AIDS should periodically receive continuing education about HIV and other STD infections to assure they have the most current information about how widespread HIV and other STDs are, effective prevention and health education intervention strategies, and priority populations identified as most at-risk for HIV and other STD infection (14,47,48). Effective implementation of school health education is linked directly to adequate teacher training programs (49). School health education designed to decrease students' participation in risk behaviors requires that teachers have appropriate training to develop and implement school health education curricula (49). Staff development activities for health education teachers need to focus on teaching strategies that both actively engage students and facilitate their mastery of critical health information and skills (50).

Question 17: During the past two years, did you receive professional development (such as workshops, conferences, continuing education, or any other kind of in-service) on each of the following topics?

Question 18: Would you like to receive professional development on each of these health education topics?

Question 19: During the past two years, did you receive professional development (such as workshops, conferences, continuing education, or any other kind of in-service) on each of the following topics?

Question 20: Would you like to receive professional development on each of these topics?

Rationale: These questions address the importance of professional development for teachers. It is vitally important that teachers be well prepared when they begin teaching and that they continue to improve their knowledge and skills throughout their careers.(51). Districts that have made improvements in their professional development activities have seen a rise in student achievement (52,53). Staff development is associated with increased teaching of important health education topics (54). The Institute of Medicine's Committee on Comprehensive School Health Programs in Grades K-12 recommended that health education teachers should be expected to participate in ongoing, discipline specific in-service programs in order to stay abreast of new developments in their field (1).

PROFESSIONAL PREPARATION

Question 21: What was the major emphasis of your professional preparation?

Question 22: Currently, are you certified, licensed, or endorsed by the state to teach health education in middle school or high school?

Question 23: Including this school year, how many years of experience do you have teaching health education classes or topics?

Rationale: These questions measure the extent to which lead health education teachers are formally trained in the topic of health education as well as the teaching experience and credentials of the lead health education teacher. Health education teachers need to be academically prepared and specifically qualified on the subject of health (50). In addition, pre-service training in health education is associated with increased teaching of important health education topics (54).

REFERENCES

1. Institute of Medicine. *Schools and Health: Our Nation's Investment*. Washington, DC: National Academy Press, 1997.
2. Lohmann DK, Wooley SF. *Comprehensive School Health Education*. In: Marx E, Wooley SF, eds. *Health is Academic*. New York: Teachers College Press, 1998:43–66.
3. The Joint Committee on National Health Education Standards. *National Health Education Standards: Achieving Excellence (2nd Edition)*. Atlanta: American Cancer Society, 2007.
4. U.S. Department of Health and Human Services. *Healthy People 2010: Understanding and Improving Health* 2nd edition. Washington, DC: U.S. Government Printing Office, November 2000.
5. U.S. Department of Health and Human Services. *Preventing Tobacco Use Among Young People: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 1994.
6. U.S. Department of Health and Human Services. *Reducing Tobacco Use: A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Public Health Service, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2000.
7. Sussman S. School-based tobacco use prevention and cessation: where are we going? *American Journal of Health Behavior* 2001; 25(3):191-9.
8. Dent CW, Sussman S, Stacy AW, Craig S, Burton D, Flay BR. Two-year behavior outcomes of project towards no tobacco use. *Journal of Consulting and Clinical Psychology* 1995; 63(4): 676-677.
9. Botvin GJ, Baker E, Dusenbury L, Botvin EM, Diaz T. Long-term follow-up results of a randomized drug abuse prevention trial in a White middle-class population. *Journal of the American Medical Association* 1995; 273(14):1106-1112.
10. Lantz PM, Jacobson PD, Warner KE, Wasserman J, Pollack HA, Berson J, Ahlstrom A. Investing in youth tobacco control: a review of smoking prevention and control strategies. *Tobacco Control* 2000; 9:47-63.
11. Rooney BL, Murray DM. A meta-analysis of smoking prevention programs after adjustment for errors in the unit of analysis. *Health Education Quarterly* 1996; 23(1):48-64.
12. Bruvold WH. A meta-analysis of adolescent smoking prevention programs. *American Journal of Public Health* 1993; 83(6):872-80.
13. Coyle SL, Boruch RF, Turner CF (Eds.). *Evaluating AIDS prevention programs* (Expanded Ed.). Washington, DC: National Academy Press, 1991.
14. Kirby D, Laris BA, Rolleri L. *Sex and HIV education programs for youth: Their impact and important characteristics*. Washington DC: Family Health International, 2006. Available at www.etr.org/recapp/programs/SexHIVedProgs.pdf.
15. Pedlow CT, Carey MP. Developmentally appropriate sexual risk reduction interventions for adolescents: rationale, review of interventions, and recommendations for research and practice. *Annals of Behavioral Medicine* 2004; 27(3):172-184.
16. Kirby D, Rolleri L, Wilson MM. *Tool to Assess the Characteristics of Effective Sex and STD/HIV Education Programs*. Washington, DC: Healthy Teen Network, 2007.

17. Food and Nutrition Board, Institute of Medicine, Committee on Prevention of Obesity of Children and Youth, Schools. In: JP Koplan, CT Liverman and VI Kraak, Editors, *Preventing Childhood Obesity: Health in the Balance*. National Academy Press, Washington, DC, 2005, pp. 237–284.
18. American Dietetic Association, Society for Nutrition Education, and American School Food Service Association. Nutrition services: an essential component of comprehensive school health programs. *Journal of Nutrition Education and Behavior* 2003; 35(2):57-67.
19. Ralston K, Buzby J, Guthrie J. *A Healthy School Meal Environment*. United States Department of Agriculture, Economic Research Service, Food Assistance and Nutrition Research Report Number 34-5. Available at: <http://www.ers.usda.gov/publications/fanrr34/fanrr34-5/fanrr34-5.pdf>.
20. US Department of Agriculture. *Changing the Scene: Improving the School Nutrition Environment*. Available at: <http://www.fns.usda.gov/tn/Resources/changing.html>.
21. CDC. Guidelines for school health programs to promote lifelong healthy eating. *MMWR* 1996; 45(No. RR-9).
22. CDC. School health index. Atlanta, GA: US Department of Health and Human Services, CDC, 2006. Available at <http://www.cdc.gov/healthyyouth/shi>
23. Hoelscher D, Feldman H, Johnson C, et al. School-based health education programs can be maintained over time: results from the CATCH institutionalization study. *Preventive Medicine* 2004; 38(5):594-606.
24. Pate RR, Davis MG, Robinson TN, Stone EJ, McKenzie TL, Young JC. Promoting physical activity in children and youth: a leadership role for schools. *Circulation* 2006; 114:1-11.
25. Pate R, Pratt M, Blair SN et al. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *Journal of the American Medical Association* 1995; 273(5):402-7.
26. CDC. *HIV Prevention in the Third Decade*. Atlanta, GA: US Department of Health and Human Services, CDC; 2003. Available at http://www.cdc.gov/hiv/HIV_3rdDecade/. 27. CDC. *HIV/AIDS Surveillance Report, 2004*. Vol. 16. Atlanta, GA: US Department of Health and Human Services, CDC; 2005:1–46.
28. Anderson RN, Smith BL. Deaths: leading causes for 2002. *National Vital Statistics Reports* 2005; 53(17):67–70.
29. CDC. Diagnosis of HIV/AIDS—32 states, 2000-2003. *MMWR* 2004 53:1106-1110.
30. CDC. National Center for HIV, STD and TB Prevention, Divisions of HIV/AIDS Prevention. *HIV/AIDS among Hispanics, 2007*. Available at <http://www.cdc.gov/hiv/pubs/facts/hispanic.htm>.
31. Eaton DK, Kann L, Kinchen S, et al. Youth risk behavior surveillance—United States, 2005. *MMWR* 2006; 55(No. SS-5).
32. Gilliland L, Scully, J.. STI-HIV Prevention: A Model Program in a School-Based Health Center. *Nursing Clinics of North America* 2005; 40:681-688.
33. Ginsburg KR, Slap GB, Cnaan A, et al.. Adolescents' perceptions of factors affecting their decisions to seek future healthcare. *Journal of the American Medical Association* 1995; 273:1913-8.
34. Ford CA, Millstein SG, Halpern-Felsher BL, Irwin CE. Influence of physician confidentiality assurances on adolescents' willingness to disclose information and seek future healthcare. *Journal of the American Medical Association* 1997; 278:1029-34.
35. Lieu T, Newacheck P, McManus M. Race, ethnicity, and access to ambulatory care among US adolescents. *American Journal of Public Health* 1993; 83:960-5.
36. Allensworth D, Kolbe L. The comprehensive school health program: state of the art. *Journal of School Health* 1987; 63:14–20.
37. Golan M, Crow S. Targeting parents exclusively in the treatment of childhood obesity: long term results. *Obesity Research* 2004;12:357-361.
38. Epstein LH, Voloski A, Wing RR, McCurley J. Ten-year outcomes of behavioral family based treatment for childhood obesity. *Health Psychology* 1994;13:373-83.
39. Splett PL, Erickson CD, Belseth SB, Jensen C. Evaluation and sustainability of the healthy learners asthma initiative. *Journal of School Health* 2006; 76(6):276-282.
40. Erickson, CD, Splett PL, Mullett SS, Jensen C, Belseth SB. The healthy learner model for student chronic condition management-Part II: The Asthma Initiative." *Journal of School Nursing* 2006; 22(6):319-329.
41. Levy M., Heffner B, Stewart T, Beeman G. The efficacy of asthma case management in an urban school district in reducing school absences and hospitalizations for asthma. *Journal of School Health* 2006; 76(6):320-324.
42. Szapocznik J, Coatsworth JD. An ecodevelopmental framework for organizing risk and protection for drug abuse: A developmental model of risk and protection. In M Glantz & CR Hartel (Eds.), *Drug Abuse: Origins and Interventions* (pp. 331–366). Washington, DC: American Psychological Association, 1999.
43. Pequegnat W, Szapocznik J. The role of families in preventing and adapting to HIV/AIDS: Issues and answers. In W Pequegnat & J Szapocznik (Eds.), *Working with families in the era of HIV/AIDS*. Thousand Oaks, CA: Sage Publications, 2000.
44. Nader PR, Sellers De, Johnson CC, Perry CL, Stone EJ, Cook KC, Bebchuk J, Luepker RV. The effect of adult participation in a school-based family intervention to improve children's diet and physical activity: the Child and Adolescent Trial for Cardiovascular Health. *Preventive Medicine* 1996;. 25:455-464.
45. Perry CL, Luepker RV, Murray DM, Kurth C, Mullis R, Crockett S, Jacobs DR. Parent involvement with children's health promotion: the Minnesota Home Team. *American Journal of Public Health* 1988; 78(9):1156-1160.
46. Hopper CA, Gruber MB, Munoz KD, Herb RA. Effect of including parents in a school-based exercise and nutrition program for children. *Research Quarterly for Exercise and Sport*. 1992; 63(3):315-321.
47. Center for AIDS Prevention Studies (CAPS) and the AIDS Research Institute, University of California, San Francisco. *What Works Best in Sex/HIV Education?* San Francisco: University of California San Francisco, 2006.
48. Council of Chief State School Officers. *What Education Leaders Should Know About Forming Partnerships to Prevent Sexual-Risk Behaviors in School-Aged Youth*. Washington, DC: Author, 2005.

49. Allensworth, D. Health education: state of the art. *Journal of School Health* 1993; 63:14–20.
50. National Commission on the Role of the School and the Community to Improve Adolescent Health. *Code Blue: Uniting for Healthier Youth*. Alexandria, VA: National Association of State Boards of Education, 1990.
51. Public Education Network. *Teacher Professional Development: A Primer for Parents and Community Members*. Washington, DC: Public Education Network, 2004.
52. Togneri W, Anderson SE. *Beyond Islands of Excellence: What Districts Can Do to Improve Instruction and Achievement in all Schools*. Washington, DC: Learning First Alliance, 2003.
53. Miles KH, Darling-Hammond L. *Rethinking the Allocation of Teaching Resources: Some Lessons From High-Performing Schools*. Philadelphia: Consortium for Policy Research in Education, 1997.
54. Jones SE, Brener ND, McManus T. The relationship between staff development and health instruction in schools in the United States. *American Journal of Health Education* 2004; 35:2-10.